

This article was downloaded by: [59.63.10.160]

On: 18 March 2015, At: 17:53

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Communicative & Integrative Biology

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/kcib20>

### Observational natural history and morphological taxonomy are indispensable for future challenges in biodiversity and conservation

Michael Staab<sup>a</sup>, Michael Ohl<sup>b</sup>, Chao-Dong Zhu<sup>c</sup> & Alexandra-Maria Klein<sup>a</sup>

<sup>a</sup> Chair of Nature Conservation and Landscape Ecology; Institute of Earth and Environmental Sciences; University of Freiburg; Freiburg, Germany

<sup>b</sup> Museum für Naturkunde; Leibniz Institute for Research on Evolution and Biodiversity; Berlin, Germany

<sup>c</sup> Key Laboratory of Zoological Systematics and Evolution; Institute of Zoology; Chinese Academy of Sciences; Beijing, PR China

Accepted author version posted online: 04 Mar 2015.



CrossMark

[Click for updates](#)

To cite this article: Michael Staab, Michael Ohl, Chao-Dong Zhu & Alexandra-Maria Klein (2015) Observational natural history and morphological taxonomy are indispensable for future challenges in biodiversity and conservation, *Communicative & Integrative Biology*, 8:1, e992745, DOI: [10.4161/19420889.2014.992745](https://doi.org/10.4161/19420889.2014.992745)

To link to this article: <http://dx.doi.org/10.4161/19420889.2014.992745>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Versions of published Taylor & Francis and Routledge Open articles and Taylor & Francis and Routledge Open Select articles posted to institutional or subject repositories or any other third-party website are without warranty from Taylor & Francis of any kind, either expressed or implied, including, but not limited to, warranties of merchantability, fitness for a particular purpose, or non-infringement. Any opinions and views expressed in this article are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor & Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

It is essential that you check the license status of any given Open and Open Select article to confirm conditions of access and use.

# Observational natural history and morphological taxonomy are indispensable for future challenges in biodiversity and conservation

Michael Staab<sup>1,\*</sup>, Michael Ohl<sup>2</sup>, Chao-Dong Zhu<sup>3</sup>, and Alexandra-Maria Klein<sup>1</sup>

<sup>1</sup>Chair of Nature Conservation and Landscape Ecology; Institute of Earth and Environmental Sciences; University of Freiburg; Freiburg, Germany; <sup>2</sup>Museum für Naturkunde; Leibniz Institute for Research on Evolution and Biodiversity; Berlin, Germany; <sup>3</sup>Key Laboratory of Zoological Systematics and Evolution; Institute of Zoology; Chinese Academy of Sciences; Beijing, PR China

**G**lobal biodiversity is rapidly declining, leading inevitably to a loss of ecosystem functionality when species and their associated life-history traits vanish. Unfortunately, even in the 21<sup>st</sup> century, a large proportion of Earth's species are yet unknown and also for most described species science lacks a deeper understanding of the functional role of species and thus of ecosystems. In this Addendum we use the recent discovery of a new spider wasp with a unique natural history as an example to emphasize the importance to conduct basic observational natural history and traditional taxonomic research. We aim to encourage such 'old-fashioned' research and biologists from various research fields to report the many fascinating phenomena holding valuable natural history information they may encounter. Such detailed knowledge on species, their life-history traits, and their trophic interactions will be crucial to reliably address the challenges global change brings to the persistence of ecosystems.

Biodiversity on Earth is facing numerous threats, most prominently by a large-scale exploitation of natural habitats to fulfill the resource demands of an exponentially growing human population.<sup>1</sup> Species extinction rates have reached alarming levels and this situation is predicted to persist throughout the 21<sup>st</sup> century.<sup>2,3</sup> As ecosystem functioning is closely intertwined with biodiversity,<sup>4,5</sup> the impact of such extinction rates are even more dreadful. While species extinctions should be avoided for ethical reasons alone, the loss of species also increases the risk of losing morphological and

behavioral trait characteristics of communities, thus affecting the functionality of ecosystems.

Unfortunately, the most endangered ecosystems which are under the highest land-use pressure harbor globally the highest number of undescribed species.<sup>6,7</sup> As there is evidence that a sixth mass extinction is already under way,<sup>8</sup> it is inevitable that species have gone and will go extinct together with their unique trait characteristics, some of them even before they are known to science. Famous examples for this are the Moa, an order of large flightless birds that vanished very shortly after the arrival of the first humans on New Zealand.<sup>9</sup> However, extinction before scientific recognition is certainly not restricted to conspicuous megafauna.<sup>10</sup>

Fortunately, we are living in the age of biodiversity discovery and never before in the history of science have new species been described at higher rates.<sup>11</sup> These findings contradict the widespread claims of a current 'taxonomy crisis' (see e.g. ref. <sup>12</sup>). Despite the 'dusty' reputation<sup>13</sup> and the, in some parts of the scientific community, low valuation of natural history and taxonomy research (see ref. <sup>14</sup>), there have been approximately 20,000 new species descriptions per year since 2009, raising the number of formally described species to almost 2 million.<sup>15</sup> DNA-based approaches, including DNA taxonomy,<sup>16</sup> DNA barcoding,<sup>17</sup> and integrative taxonomy<sup>18,19</sup> complement and speed up the discovery rates of new species.

In a recent paper,<sup>20</sup> we described a new species of spider wasp which shows a striking nesting behavior: *Deuteragenia*

**Keywords:** BEF-China, ecosystem functioning, global change, integrative taxonomy, species extinctions, species interactions, trait characteristics

\*Correspondence to: Michael Staab; Email: michael.staab@nature.uni-freiburg.de

Submitted: 09/22/2014

Accepted: 09/24/2014

<http://dx.doi.org/10.4161/19420889.2014.992745>

Addendum to: Staab M, Ohl M, Zhu CD, Klein AM (2014) A unique nest-protection strategy in a new species of spider wasp. PLoS ONE 9: e101592. doi:10.1371/journal.pone.0101592.

*ossarium* Ohl, 2014, protects uniquely in the entire animal kingdom its own progeny by a nest-closing chamber filled with dead ants. This discovery was fascinating but totally unexpected; actually we were, as part of a large Chinese-German research cooperation (www.bef-china.de, see ref. <sup>21</sup>), conducting a study to disentangle the effects of tree diversity on insect communities. Doing so, we collected solitary cavity-nesting Hymenoptera with trap nests,<sup>22</sup> in which we found the new species among several other, mostly described bee and wasp species. Thus we see our study as an example for the importance to report on-the-side observations that may frequently occur during larger studies. We urge biologists, in particular graduate students who usually do most of the basic research, to share and publish the intriguing phenomena they may encounter. Otherwise many fascinating aspects of species' biology will never get publicly known.

We would also like to highlight that *D. ossarium* and its unique life-history were only discovered by conducting a detailed observational study. Even in the 21st century, in the time of high-technology science, for most plant and animal species barely more than the name and the type locality is known, showing the urgent need for basic natural history studies. Moreover, many new species are described without knowing any associated information and most of them only decades after the initial collection.<sup>23</sup> In our case, crucial aspects of the life-history of *D. ossarium*, namely the very low susceptibility to parasitism and that the species, despite being a spider-hunting wasp, is possibly a significant predator of the most abundant ant species in its range, would have stayed unnoticed if we had only detected the species by molecular methods or if we had only described the species morphologically. Our discovery demonstrates that molecular methods to identify new species can, if part of an integrative holistic taxonomic approach,<sup>18,19</sup> complement but not displace classical observations, as the complete natural history of a species will only be caught by detailed observational and experimental research.<sup>13,24</sup>

We realize and understand that, due to the limited availability of funding and

manpower, it will only for a minority of species be possible to conduct natural history studies. For most species, the life-history, the functional roles in an ecosystem, and the trophic interactions will likely ever stay unknown or obscure. This is particularly true for invertebrates, which account for the largest part of animal biodiversity and biomass<sup>25</sup> and which are indispensable for ecosystem functioning and services.<sup>26,27</sup> We are convinced that there are very many functionally important life-history traits yet unknown across the tree of life, both in already known species and in species awaiting discovery and description. Knowledge on these species and their traits will be necessary to reliably assess the functionality of single species and consequently of the ecosystems the species live in. Accurate and holistic data on species, traits, and communities will be vitally important to predict responses of species and ecosystems to global change,<sup>28</sup> to develop effective conservation schemes, and to preserve critical ecosystem functions that depend to a high degree on biodiversity.<sup>4,5</sup> We emphasize that our proposal for taxonomy and natural history studies is not restricted to pristine habitats. Our discovery<sup>20</sup> has been made in South-East China, a region where the original species-rich subtropical forests are under high land-use pressure and have largely been converted to agricultural land or forestry monocultures.<sup>29,30</sup> Thus it is very likely that in this region several species have gone extinct together with their unique life history, behavior and morphology long before they were observed, recognized or analyzed.

#### Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

#### Acknowledgments

We thank the Editor in Chief, František Baluška, for his invitation and encouragement to write this addendum. We also thank Tamar Marcus and David Walmsley for stimulating discussions and for proof-reading an earlier version of the manuscript. The original work on which this addendum is based on was part of the

'Biodiversity-Ecosystem Functioning China Project' (BEF-China) funded by the German Research Foundation (DFG; grants FOR 891/2, KL 1849/6-1).

#### References

1. McKee JK, Sciulli PW, Foose CD, Waite TA. Forecasting global biodiversity threats associated with human population growth. *Biol Conserv* 2004; 115:161-4; [http://dx.doi.org/10.1016/S0006-3207\(03\)00099-5](http://dx.doi.org/10.1016/S0006-3207(03)00099-5)
2. Butchart SHM, Walpole M, Collen B, van Strien A, Scharlemann JPW, Almond REA, Baillie JEM, Bomhard B, Brown C, Bruno J, et al. Global biodiversity: Indicators of recent declines. *Science* 2010; 328: 1164-8; PMID:20430971; <http://dx.doi.org/10.1126/science.1187512>
3. Pereira HM, Leadley PW, Proenca V, Alkemade R, Scharlemann JPW, Fernandez-Manjarres JF, Araujo MB, Balvanera P, Biggs R, Cheung WWL, et al. Scenarios for global biodiversity in the 21st century. *Science* 2010; 330:1496-501; PMID:20978282; <http://dx.doi.org/10.1126/science.1196624>
4. Balvanera P, Pfisterer AB, Buchmann N, He JS, Nakashizuka T, Raffaelli D, Schmid B. Quantifying the evidence for biodiversity effects on ecosystem functioning and services. *Ecol Lett* 2006; 9:1146-56; PMID:16972878; <http://dx.doi.org/10.1111/j.1461-0248.2006.00963.x>
5. Cardinale BJ, Duffy JE, Gonzalez A, Hooper DU, Perings C, Venail P, Narwani A, Mace GM, Tilman D, Wardle DA, et al. Biodiversity loss and its impact on humanity. *Nature* 2012; 486:59-67; PMID:22678280; <http://dx.doi.org/10.1038/nature11148>
6. Joppa LN, Roberts DL, Myers N, Pimm SL. Biodiversity hotspots house most undiscovered plant species. *Proc Natl Acad Sci U S A* 2011; 108:13171-6; PMID:21730155; <http://dx.doi.org/10.1073/pnas.1109389108>
7. Guénard B, Weiser MD, Dunn RR. Global models of ant diversity suggest regions where new discoveries are most likely are under disproportionate deforestation threat. *Proc Natl Acad Sci U S A* 2012; 109:7368-73; PMID:22529355; <http://dx.doi.org/10.1073/pnas.1113867109>
8. Barnosky AD, Matzke N, Tomiya S, Wogan GOU, Swartz B, Quental TB, Marshall C, McGuire JL, Lindsey EL, Maguire KC, et al. Has the Earth's sixth mass extinction already arrived? *Nature* 2011; 471:51-7; PMID:21368823; <http://dx.doi.org/10.1038/nature09678>
9. Holdaway RN, Jacomb C. Rapid extinction of the Moas (Aves: Dinorhithiformes): Model, test, and implications. *Science* 2000; 287:2250-4; PMID:10731144; <http://dx.doi.org/10.1126/science.287.5461.2250>
10. Richling I, Boucher P. Extinct even before scientific recognition: A remarkable radiation of helicidid snails (Helicinidae) on the Gambier Islands, French Polynesia. *Biodivers Conserv* 2013; 22:2433-68; <http://dx.doi.org/10.1007/s10531-013-0496-2>
11. Costello MJ, May RM, Stork NE. Can we name Earth's species before they go extinct? *Science* 2013; 339:413-6; PMID:23349283; <http://dx.doi.org/10.1126/science.1230318>
12. Agnarsson I, Kuntner M. Taxonomy in a changing world: Seeking solutions for a science in crisis. *Syst Biol* 2007; 56:531-9; PMID:17562477; <http://dx.doi.org/10.1080/10635150701424546>
13. Ebach MC, Holdrege C. More taxonomy, not DNA barcoding. *Bio Sci* 2005; 55:822-3; [http://dx.doi.org/10.1641/0006-3568\(2005\)055\[0823:MTNDBJ\]2.0.CO;2](http://dx.doi.org/10.1641/0006-3568(2005)055[0823:MTNDBJ]2.0.CO;2)
14. Wägele H, Klusmann-Kolb A, Kuhlmann M, Haszprunar G, Lindberg D, Koch A, Wägele JW. The taxonomist - an endangered race. A practical proposal for

- its survival. *Front Zool* 2011; 8:25; PMID:22029904; <http://dx.doi.org/10.1186/1742-9994-8-25>
15. IISE. State of observed species. Tempe, AZ, U S A: International Institute for Species Exploration 2011; retrieved 22.09.2014; <http://timgostony.com/iise/temp/>
  16. Tautz D, Arctander P, Minelli A, Thomas RH, Vogler AP. A plea for DNA taxonomy. *Trends Ecol Evol* 2003; 18:70-4; [http://dx.doi.org/10.1016/S0169-5347\(02\)00041-1](http://dx.doi.org/10.1016/S0169-5347(02)00041-1)
  17. Hebert PDN, Cywinska A, Ball SL, DeWaard JR. Biological identifications through DNA barcodes. *Proc R Soc B* 2003; 270:313-21; PMID:12614582; <http://dx.doi.org/10.1098/rspb.2002.2218>
  18. Dayrat B. Towards integrative taxonomy. *Biol J Linn Soc* 2005; 85:407-15; <http://dx.doi.org/10.1111/j.1095-8312.2005.00503.x>
  19. Challenges for Taxonomic Research Working Group. Challenges for taxonomic research in the age of 'omics' technologies. Halle/Saale, Germany: Leopoldina Nationale Akademie der Wissenschaften, 2014; retrieved 22.09.2014; <http://www.leopoldina.org/en/policy-advice/working-groups/completed-working-groups/challenges-for-taxonomic-research/>
  20. Staab M, Ohl M, Zhu CD, Klein AM. A unique nest-protection strategy in a new species of spider wasp. *PLoS One* 2014; 9:e101592; PMID:24987876; <http://dx.doi.org/10.1371/journal.pone.0101592>
  21. Bruehlheide H, Böhnke M, Both S, Fang T, Assmann T, Baruffol M, Bauhus J, Buscot F, Chen XY, Ding BY, et al. Community assembly during secondary forest succession in a Chinese subtropical forest. *Ecol Monogr* 2011; 81:25-41; <http://dx.doi.org/10.1890/09-2172.1>
  22. Tscharnkte T, Gathmann A, Steffan-Dewenter I. Bioindication using trap-nesting bees and wasps and their natural enemies: community structure and interactions. *J Appl Ecol* 1998; 35:708-19; <http://dx.doi.org/10.1046/j.1365-2664.1998.355343.x>
  23. Fontaine B, Perrard A, Bouchet P. 21 years of shelf life between discovery and description of new species. *Curr Biol* 2012; 22:943-4; PMID:23174292; <http://dx.doi.org/10.1016/j.cub.2012.10.029>
  24. Ebach MC, Holdrege C. DNA barcoding is no substitute for taxonomy. *Nature* 2005; 434:697; PMID:15815602; <http://dx.doi.org/10.1038/434697b>
  25. Wilson EO. The little things that run the world. *Conserv Biol* 1987; 1:344-6; <http://dx.doi.org/10.1111/j.1523-1739.1987.tb00055.x>
  26. Folgarait PJ. Ant biodiversity and its relationship to ecosystem functioning: a review. *Biodivers Conserv* 1998; 7:1221-44; <http://dx.doi.org/10.1023/A:1008891901953>
  27. Losey JE, Vaughan M. The economic value of ecological services provided by insects. *Bio Sci* 2006; 56:311-23; [http://dx.doi.org/10.1641/0006-3568\(2006\)56\[311:revoes\]2.0.co;2](http://dx.doi.org/10.1641/0006-3568(2006)56[311:revoes]2.0.co;2)
  28. Cahill AE, Aiello-Lammens ME, Fisher-Reid MC, Hua X, Karanewsky CJ, Ryu HY, Sbeglia GC, Spagnolo F, Waldron JB, Warsi O, et al. How does climate change cause extinction? *Proc R Soc B* 2013; 280:20121890; PMID:23075836; <http://dx.doi.org/10.1098/rspb.2012.1890>
  29. López-Pujol J, Zhang FM, Ge S. Plant biodiversity in China: Richly varied, endangered, and in need of conservation. *Biodivers Conserv* 2006; 15:3983-4026; <http://dx.doi.org/10.1007/s10531-005-3015-2>
  30. Hansen MC, Potapov PV, Moore R, Hancher M, Turubanova SA, Tyukavina A, Thau D, Stehman SV, Goetz SJ, Loveland TR, et al. High-resolution global maps of 21st-century forest cover change. *Science* 2013; 342:850-3; PMID:24233722; <http://dx.doi.org/10.1126/science.1244693>